



Short Communication

An Examination of the Indirect Effect of Anxiety Sensitivity in terms of Asthma and Smoking Cessation Processes



Alison C. McLeish^{a,*}, Samantha G. Farris^b, Adrienne L. Johnson^a, Jonathan A. Bernstein^c, Michael J. Zvolensky^{b,d}

^a Department of Psychology, University of Cincinnati, PO Box 2120376, Cincinnati, OH, 45221–0376, USA

^b Department of Psychology, University of Houston, 126 Heyne Building, Houston, TX, 77204–5022, USA

^c Department of Internal Medicine, Division of Immunology, Allergy, and Rheumatology, University of Cincinnati College of Medicine, 3225 Eden Avenue, Cincinnati, OH, 45267–0563, USA

^d The University of Texas MD Anderson Cancer Center, 1515 Holcombe Blvd., Houston, TX, 77030, USA

HIGHLIGHTS

- Smoking is more common among individuals with asthma compare to those without
- Anxiety sensitivity negative impacts both smoking and asthma
- Examined the indirect effect of anxiety sensitivity in terms of asthma and cessation processes
- Significant indirect effect for nicotine dependence motives and motivation to quit
- No significant indirect effect for duration of longest quit attempt.

ARTICLE INFO

Article history:

Received 30 January 2015

Received in revised form 7 April 2015

Accepted 16 June 2015

Available online 23 June 2015

Keywords:

anxiety sensitivity
asthma
tobacco
motives
smoking cessation

ABSTRACT

Introduction: Despite the increased rates of smoking and poor cessation outcomes among individuals with asthma relative to those without, little scholarly attention has examined mechanisms linking asthma to smoking cessation processes. The current study sought to examine the indirect effect of anxiety sensitivity in terms of asthma and smoking cessation processes (i.e., duration of longest quit attempt, motivation to quit smoking, smoking dependence motives).

Methods: Participants were 90 regular daily smokers: 43 with asthma (51.2% male, $M_{age} = 38.0$ years, $SD = 12.5$) and 47 without asthma (46.8% male, $M_{age} = 35.4$ years, $SD = 11.2$) who were participating in a larger smoking cessation study. Data from the baseline (pre-quit attempt) assessment session were used.

Results: After accounting for the effects of gender, race, daily smoking rate, and negative affectivity, asthma status was indirectly related to motives for smoking related to nicotine dependence and motivation to quit smoking through anxiety sensitivity. There was no significant indirect effect for duration of longest quit attempt.

Conclusions: These findings suggest that smokers with asthma may be particularly fearful of physiological arousal, which in turn, may account for greater motivation to quit smoking, but also stronger dependence motives for smoking.

© 2015 Elsevier Ltd. All rights reserved.

1. Introduction

Cigarette smoking is more common among individuals with asthma compared to those without (Gwynn, 2004; McLeish, Cogle, & Zvolensky, 2011), resulting in greater asthma severity, poorer asthma control, and more frequent healthcare utilization (Althuis, Sexton, & Prybylski, 1999; Eisner & Iribarren, 2007; McLeish & Zvolensky, 2010; Siroux, Pin, Oryszczyn, Le Moual, & Kauffmann, 2000). Although quitting smoking results in significant improvements in lung function, reductions in asthma medication use, and improved quality of life (Chaudhuri et al.,

2006; Tønnesen et al., 2005), little is known about factors that influence smoking cessation among smokers with asthma. This lack of attention is unfortunate as smokers with asthma are less successful than smokers without when trying to quit, despite making more quit attempts (Avallone et al., 2013; Fennerty et al., 1987, 1987; Tønnesen et al., 2005).

One factor that may be important to examine in terms of asthma-smoking relations is the cognitive risk factor of anxiety sensitivity, defined as the fear of arousal-related sensations (McNally, 2002; Reiss, Peterson, Gursky, & McNally, 1986). Research suggests that anxiety sensitivity serves as a risk factor for both smoking and asthma, resulting in poorer outcomes in both domains. For example, anxiety sensitivity predicts panic in response to asthma symptoms, poorer asthma control, and poorer asthma related quality of life (Avallone, McLeish, Luberto, &

* Corresponding author. Tel.: +1 513 556 5123; fax: +1 513 556 1904.
E-mail address: alison.mcleish@uc.edu (A.C. McLeish).

Bernstein, 2012; Carr, Lehrer, & Hochron, 1995; McLeish, Zvolensky, & Luberto, 2011) as well as greater perceived barriers to cessation and increased withdrawal symptoms and risk of lapse and relapse during a quit attempt (Johnson, Farris, Schmidt, Smits, & Zvolensky, 2013; Johnson, Stewart, Rosenfield, Steeves, & Zvolensky, 2012; Mullane et al., 2008).

In the only study, to date, examining the role of anxiety sensitivity among smokers with asthma, McLeish, Johnson, Avallone, and Zvolensky (in press) found that greater anxiety sensitivity significantly predicted greater barriers to cessation and reasons for quitting related to health concerns and self-control. These findings suggest that anxiety sensitivity does indeed relate to important smoking cessation processes among smokers with asthma; however, this study only examined the main effect of anxiety sensitivity, asthma diagnosis was not objectively verified, and there was no comparison to smokers without asthma. Therefore, the current study sought to build on past work by examining the indirect effect of anxiety sensitivity in terms of asthma status and smoking cessation processes in a sample of smokers with and without asthma enrolled in a smoking cessation study. It was hypothesized that there would be a significant indirect effect of asthma diagnosis on shorter duration of longest quit attempt, greater smoking cessation motives, and greater smoking dependence motives through anxiety sensitivity.

2. Method

2.1. Participants

Participants were 90 daily cigarette smokers between the ages of 18 and 65. For inclusion in the study, participants had to: (1) be a regular smoker (≥ 10 cigarettes per day) for at least one year; (2) have expired carbon monoxide (CO) levels ≥ 8 parts per million (ppm); (3) report a motivation to quit smoking of at least 5 out of 10; and (4) be interested in quitting smoking in the next month. Participants were excluded from the study based on: (1) current substance dependence (excluding nicotine dependence); (2) decreased smoking rate by more than a half in the past six months; or (3) current, regular use of other tobacco products. Participants in the smokers with asthma group were required to have received a physician diagnosis of asthma prior to the onset of smoking and meet criteria for asthma diagnosis verification using spirometry. The smokers without asthma group ($n = 47$; 46.8% male, $M_{age} = 35.4$ years, $SD = 11.2$) was 56.5% Caucasian, 39.1% African American, and 4.3% Other. On average, smokers without asthma smoked 17.2 ($SD = 11.1$) cigarettes per day and had been regular smokers for 16.2 years ($SD = 9.8$). The smokers with asthma group ($n = 43$; 51.2% male, $M_{age} = 38.0$ years, $SD = 12.5$) was 39.5% Caucasian, 58.1% African American, and 2.3% Other. On average, smokers with asthma smoked 22.3 ($SD = 22.0$) cigarettes per day and had been regular smokers for 19.9 years ($SD = 11.9$).

2.2. Measures

2.2.1. Asthma diagnosis

Asthma status was verified by spirometry using a KoKo Legend spirometer (nSpire Health, Inc., Longmont, CO). Individuals who demonstrated a reduction in FEV₁ and FEV₁/FVC values relative to predicted values, with 12% or greater improvement after administration of short-acting bronchodilator (or 20% or greater improvement in FEF_{25–75}) were considered to have a positive asthma status (Alberts, Ferris, Brooks, & Goldman, 1994; National Heart Lung & Blood Institute, 2007).

2.2.2. Expired carbon monoxide (CO)

Biochemical verification of smoking status was completed by CO analysis of breath samples using a Bedfont Micro 4 Smokerlyzer CO

Monitor (coVita, Haddonfield, NJ). Obtained values ≥ 8 ppm were considered indicative of regular smoking (Benowitz et al., 2002).

2.2.3. Smoking History Questionnaire (SHQ).

Smoking history and pattern was assessed with the SHQ (Brown, Lejuez, Kahler, & Strong, 2002). The SHQ includes items pertaining to smoking rate, years of being a daily smoker, and duration of longest quit attempt in days.

2.2.4. Positive Affect Negative Affect Schedule (PANAS).

The PANAS (Watson, Clark, & Tellegen, 1988) is a measure of general positive and negative emotional states commonly used in psychopathology research (Watson, 2000). In the present study, only the negative affect subscale (PANAS-NA) was used.

2.2.5. Anxiety Sensitivity Index-3 (ASI-3)

The ASI-3 (Taylor et al., 2007) is an 18-item self-report measure that assesses the degree to which participants fear negative consequences stemming from anxiety symptoms. The ASI-3 has demonstrated the strongest psychometric properties of any current measure of AS (Taylor et al., 2007).

2.2.6. Motivational Aspects of Smoking Cessation (MASC).

The MASC is an 11-item self-report measure that assesses the degree to which participants are motivated to quit smoking (Rundmo, Smedslund, & Götestam, 1997). The MASC has demonstrated good internal consistency and validity (Rundmo et al., 1997).

2.2.7. Brief Wisconsin Inventory for Smoking Dependence Motives (WISDM-37)

The WISDM-37 (Smith et al., 2010) is 37-item self-report measure of tobacco dependence. It is a shortened version of the original 68-item Wisconsin Inventory for Smoking Dependence Motives (WISDM-68; Piper et al., 2004). The WISDM-37 consists of 11 subscales that load onto two higher order factors: (1) primary dependence motives (automaticity, loss of control, craving, tolerance) and (2) secondary dependence motives (affiliative attachment, cognitive enhancement, cue exposure/associative processes, social/environmental goals, taste, weight control, affective enhancement). The WISDM-37 shows good internal consistency, reliability, concurrent, and predictive validity (Smith et al., 2010).

2.3. Procedure

Participants were daily smokers participating in a larger study examining differences between smokers with and without asthma during a self-guided quit attempt. The current study uses data from the study's baseline assessment session, which took place prior to the quit attempt. Individuals interested in participating in the study were screened by phone and eligible participants were scheduled for an in-person baseline visit. After providing informed, written consent, participants had their smoking status and asthma status verified, and completed a self-report assessment battery. Upon completion of the visit, participants were compensated \$50.

3. Results

Model fit for the direct and indirect effects of asthma status in terms of smoking cessation motives and smoking dependence motives was very good (see Fig. 1; $X^2(26) = 30.189$, $p = .260$; CFI = .961, RMSEA = .055). Having an asthma diagnosis was not significantly associated with smoking more cigarettes per day, non-white race, or female gender. Based on modification indices, a number of changes were made to the model. Race was allowed to correlate with smoking rate ($r = .03$, $p = ns$), gender ($r = .21$, $p = .041$), and anxiety sensitivity ($r = .28$, $p = .008$). Negative affectivity was retained as a correlated




Fig. 1. Path Model. Covariates and errors terms are not displayed for ease of viewing. Grey lines are non-significant direct effects; thick black regression lines indicate significant direct effects. $^*p < .05$, $^{**}p < .01$.

factor with anxiety sensitivity in the model despite the non-significant association based on theoretical relevance ($r = .03$, $p = .ns$). Gender was also allowed to correlate with negative affectivity ($r = .15$, $p = ns$). Primary and secondary smoking motives were allowed to correlate ($r = .67$, $p < .001$). Smoking rate was allowed to correlate to duration of longest quit attempt ($r = .27$, $p = .015$).

Asthma group status was significantly and positively associated with anxiety sensitivity ($\beta = .23$, $p = .031$). Asthma status was not directly related to smoking cessation motives, duration of longest quit attempt, or smoking dependence motives (grey lines in Fig. 1). Anxiety sensitivity was directly related to greater motivation to quit smoking ($\beta = .29$, $p = .005$), shorter duration of longest quit attempt ($\beta = -.22$, $p = .029$), stronger primary dependence motives ($\beta = .31$, $p = .008$), and stronger secondary dependence motives ($\beta = .40$, $p < .001$). There was a significant indirect effect of asthma group status on motivation to quit smoking through the effect of anxiety sensitivity ($a^*b = 0.99$, $SE = 0.60$ [95% CI: 0.046, 2.373]). Anxiety sensitivity also mediated the association between asthma group status and primary dependence motives ($a^*b = 0.16$, $SE = 0.09$ [95% CI: 0.010, 0.368]) and secondary dependence motives ($a^*b = 0.20$, $SE = 0.11$, [95% CI: 0.017, 0.445]). The indirect effect of asthma group status on duration of longest quit attempt through anxiety sensitivity was non-significant.

4. Discussion

As hypothesized, there was a significant indirect effect of asthma status on motivation to quit smoking, primary dependence motives, and secondary dependence motives through anxiety sensitivity. Contrary to hypothesis, there was no significant indirect effect of asthma on duration of longest quit attempt through anxiety sensitivity. These findings suggest that smokers with asthma are both more motivated to quit smoking and to smoke more compulsively and for certain situational reasons (e.g., affect regulation) due to the tendency to be fearful and misinterpret the meaning of arousal-related physical and psychological sensations. As such, anxiety sensitivity appears to serve as a mechanism underlying associations between asthma and motivational smoking cessation processes (i.e., motivation for quitting, motivations associated with nicotine dependence/problematic use). It is interesting, and perhaps unfortunate, that these are essentially competing motivations: one to stop smoking and one to continue smoking. The non-significant indirect effect for duration of longest quit attempt was unexpected; however, anxiety sensitivity appears to be most relevant early in the quit attempt (i.e., first two weeks; Zvolensky, Stewart, Vujanovic, Gavric, & Steeves, 2009), which we were unable to assess in this cross-sectional study.

There are a number of limitations to the current study that warrant consideration. First, the present cross-sectional design does not permit causal-oriented hypothesis testing. Longitudinal studies are needed to better understand how anxiety sensitivity impacts smoking cessation over time. Second, the sample, by design, consisted of regular daily smokers who smoked at least 10 cigarettes per day, limiting the generalizability of the results to individuals who smoke less than that or do not smoke on a daily basis. Finally, comorbid medical conditions were not assessed. Future studies would benefit from conducting a more comprehensive assessment of participants' medical history to ensure that findings are not due to the presence of an increased number of medical conditions among smokers with asthma. Despite these limitations, the results of this study indicate smokers with asthma may have greater difficulties quitting smoking because of their tendency to be fearful of physiological arousal and may benefit from specialized smoking cessation interventions that specifically target anxiety sensitivity.

Role of funding sources

Funding for this study was provided by American Lung Association grant SB-193355-N awarded to ACM and National Institute of Drug Abuse grant F31-DA035564 awarded to SGF. The American Lung Association and National Institute on Drug Abuse had no role in the study design, collection, analysis or interpretation of the data, writing the manuscript, or the decision to submit the paper for publication.

Contributors

ACM, MJZ, and JAB designed the study. ALJ collected the data and drafted the Abstract, Method, and References sections. SGF conducted the statistical analyses. ACM and SGF wrote the first draft of the manuscript, and ALJ, JAB, and MJZ provided significant input in revising the manuscript drafts. All authors contributed to and have approved the final manuscript.

Conflict of interest

All authors declare that they have no conflicts of interest.

References

- Alberts, W.M., Ferris, M.C., Brooks, S.M., & Goldman, A.L. (1994). The FEF25-75% and the clinical diagnosis of asthma. *Annals of Allergy*, *73*, 221–225.
- Althuis, M.D., Sexton, M., & Prybylski, D. (1999). Cigarette smoking and asthma symptom severity among adult asthmatics. *Journal of Asthma*, *36*, 257–264. <http://dx.doi.org/10.3109/02770909909075409>.
- Avallone, K.M., McLeish, A.C., Luberto, C.M., & Bernstein, J.A. (2012). Anxiety sensitivity, asthma control, and quality of life in adults with asthma. *Journal of Asthma*, *49*, 57–62. <http://dx.doi.org/10.3109/02770903.2011.641048>.
- Avallone, K.M., McLeish, A.C., Zvolensky, M.J., Kraemer, K.M., Luberto, C.M., & Jeffries, E.R. (2013). Asthma and its relation to smoking behavior and cessation motives among adult daily smokers. *Journal of Health Psychology*, *18*, 788–799. <http://dx.doi.org/10.1177/1359105312456322>.
- Benowitz, N.L., Jacob, P., Ahijevych, K., Jarvis, M.J., Hall, S., LeHouezec, J., et al. (2002). Biochemical verification of tobacco use and cessation. *Nicotine & Tobacco Research*, *4*, 149–159. <http://dx.doi.org/10.1080/14622200210123581>.

- Brown, R.A., Lejuez, C.W., Kahler, C.W., & Strong, D.R. (2002). Distress tolerance and duration of past smoking cessation attempts. *Journal of Abnormal Psychology, 111*, 180–185. <http://dx.doi.org/10.1037/0021-843X.111.1.180>.
- Carr, R.E., Lehrer, P.M., & Hochron, S.M. (1995). Predictors of panic-fear in asthma. *Health Psychology, 14*, 421–426. [http://dx.doi.org/10.1016/S0006-3223\(02\)01475-0](http://dx.doi.org/10.1016/S0006-3223(02)01475-0).
- Chaudhuri, R., Livingston, E., McMahon, A.D., Lafferty, J., Fraser, I., Spears, M., et al. (2006). Effects of smoking cessation on lung function and airway inflammation in smokers with asthma. *American Journal of Respiratory and Critical Care Medicine, 174*, 127–133. <http://dx.doi.org/10.1164/rccm.200510-1589OC>.
- Eisner, M.D., & Iribarren, C. (2007). The influence of cigarette smoking on adult asthma outcomes. *Nicotine & Tobacco Research, 9*, 53–56. <http://dx.doi.org/10.1080/14622200601078293>.
- Fennerty, A.G., Banks, J., Ebdon, P., & Bevan, C. (1987). The effect of cigarette withdrawal on asthmatics who smoke. *European Journal of Respiratory Diseases, 71*, 395–399.
- Gwynn, R. C. (2004). Risk factors for asthma in US adults: results from the 2000 behavioral risk factor surveillance system. *J. Asthma, 41*, 91–98. <http://dx.doi.org/10.1081/jas-120026066>.
- Johnson, K.A., Farris, S.G., Schmidt, N.B., Smits, J.A., & Zvolensky, M.J. (2013). Panic attack history and anxiety sensitivity in relation to cognitive-based smoking processes among treatment-seeking daily smokers. *Nicotine & Tobacco Research, 15*, 1–10. <http://dx.doi.org/10.1093/ntr/ntr332>.
- Johnson, K.A., Stewart, S., Rosenfield, D., Steeves, D., & Zvolensky, M.J. (2012). Prospective evaluation of the effects of anxiety sensitivity and state anxiety in predicting acute nicotine withdrawal symptoms during smoking cessation. *Psychology of Addictive Behaviors, 26*, 289–297. <http://dx.doi.org/10.1037/a0024133>.
- McLeish, A.C., Cogle, J.R., & Zvolensky, M.J. (2011). Asthma and cigarette smoking in a representative sample of adults. *Journal of Health Psychology, 16*, 643–652. <http://dx.doi.org/10.1177/1359105310386263>.
- McLeish, A.C., Johnson, A.L., Avallone, K.M., & Zvolensky, M.J. (2014n). *Evaluating the role of anxiety sensitivity in barriers to cessation and reasons for quitting among smokers with asthma*. (in press).
- McLeish, A.C., & Zvolensky, M.J. (2010). Asthma and cigarette smoking: A review of the empirical literature. *Journal of Asthma, 47*, 345–361. <http://dx.doi.org/10.3109/02770900903556413>.
- McLeish, A.C., Zvolensky, M.J., & Luberto, C.M. (2011). The role of anxiety sensitivity in terms of asthma control: A pilot test among young adult asthmatics. *Journal of Health Psychology, 16*, 439–444. <http://dx.doi.org/10.1177/1359105310382584>.
- McNally, R.J. (2002). Anxiety sensitivity and panic disorder. *Biological Psychiatry, 52*, 938–946. [http://dx.doi.org/10.1016/S0006-3223\(02\)01475-0](http://dx.doi.org/10.1016/S0006-3223(02)01475-0).
- Mullane, J.C., Stewart, S.H., Rhyno, E., Steeves, D., Watt, M., & Eisner, A. (2008). Anxiety, sensitivity and difficulties with smoking cessation. In A.M. Columbus (Ed.), *Advances in psychology research, Vol. 54*. (pp. 141–155). Hauppauge, NY, US: Nova Science Publishers.
- National Heart Lung and Blood Institute (2007). *National Asthma Education Program Expert Panel Report 3: Guidelines for the diagnosis and management of asthma*. (Bethesda, MD).
- Piper, M.E., Piasecki, T.M., Federman, E.B., Bolt, D.M., Smith, S.S., Fiore, M.C., et al. (2004). A multiple motives approach to tobacco dependence: the Wisconsin Inventory of Smoking Dependence Motives (WISDM-68). *Journal of Consulting and Clinical Psychology, 72*, 139–154. <http://dx.doi.org/10.1037/0022-006X.72.2.139>.
- Reiss, S., Peterson, R.A., Gursky, D.M., & McNally, R.J. (1986). Anxiety sensitivity, anxiety frequency and the prediction of fearfulness. *Behaviour Research and Therapy, 24*, 1–8. [http://dx.doi.org/10.1016/0005-7967\(86\)90143-9](http://dx.doi.org/10.1016/0005-7967(86)90143-9).
- Rundmo, T., Smedslund, G., & Götestam, K.G. (1997). Motivation for smoking cessation among the Norwegian public. *Addictive Behaviors, 22*, 377–386. [http://dx.doi.org/10.1016/S0306-4603\(96\)00056-1](http://dx.doi.org/10.1016/S0306-4603(96)00056-1).
- Siroux, V., Pin, I., Oryszczyn, M.P., Le Moual, N., & Kauffmann, F. (2000). Relationships of active smoking to asthma and asthma severity in the EGEA study. Epidemiological study on the Genetics and Environment of Asthma. *European Respiratory Journal, 15*, 470–477. <http://dx.doi.org/10.1034/j.1399-3003.2000.15.08.x>.
- Smith, S.S., Piper, M.E., Bolt, D.M., Fiore, M.C., Wetter, D.W., Cinciripini, P.M., et al. (2010). Development of the Brief Wisconsin Inventory of Smoking Dependence Motives. *Nicotine & Tobacco Research, 12*, 489–499. <http://dx.doi.org/10.1093/ntr/ntq032>.
- Taylor, S., Zvolensky, M.J., Cox, B.J., Deacon, B., Heimberg, R.G., Ledley, D.R., et al. (2007). Robust dimensions of anxiety sensitivity: development and initial validation of the Anxiety Sensitivity Index-3. *Psychological Assessment, 19*, 176–188. <http://dx.doi.org/10.1037/1040-3590.19.2.176>.
- Tønnesen, P., Pisinger, C., Hvidberg, S., Wennike, P., Bremann, L., Westin, A., et al. (2005). Effects of smoking cessation and reduction in asthmatics. *Nicotine & Tobacco Research, 7*, 139–148. <http://dx.doi.org/10.1080/14622200412331328411>.
- Watson, D. (2000). *Mood and temperament*. New York: Guilford Press.
- Watson, D., Clark, L.A., & Tellegen, A. (1988). Development and Validation of Brief Measures of Positive and Negative Affect: The PANAS Scales. *Journal of Personality and Social Psychology, 54*, 1063–1070. <http://dx.doi.org/10.1037/0022-3514.54.6.1063>.
- Zvolensky, M.J., Stewart, S.H., Vujanovic, A.A., Gavric, D., & Steeves, D. (2009). Anxiety sensitivity and anxiety and depressive symptoms in the prediction of early smoking lapse and relapse during smoking cessation treatment. *Nicotine & Tobacco Research, 11*, 323–331. <http://dx.doi.org/10.1093/ntr/ntn037>.